

Bridgepoint Plaza I, Suite 300 5918 West Courtyard Drive • Austin, TX 78730-5036 Phone 512-329-6610 • Fax 512-327-7159 • www.tmf.org

Notice of Independent Review Decision

IRO REVIEWER REPORT

DATE OF REVIEW: 03/22	2/2010
IRO CASE #:	
DESCRIPTION OF THE S	ERVICE OR SERVICES IN DISPUTE
Prescriptions for Flexeril, \	/alium, Norco and Xanax
	QUALIFICATIONS FOR EACH PHYSICIAN OR ROVIDER WHO REVIEWED THE DECISION
unrestricted license to pract	er is board certified occupational medicine with an ctice in the state of Texas. The physician is in active the treatment or proposed treatment.
REVIEW OUTCOME	
Upon independent review determination/adverse det	the reviewer finds that the previous adverse erminations should be:
⊠ Upheld	(Agree)
Overturned	(Disagree)
☐ Partially Overturned	(Agree in part/Disagree in part)
<u>-</u>	e review outcome that clearly states whether or not each of the health care services in dispute.
•	escriptions for Flexeril, Valium, Norco and Xanax are

INFORMATION PROVIDED TO THE IRO FOR REVIEW

• Information for requesting a review by an IRO – 01/29/10

- Office visit notes by Dr. 10/30/08 to 01/20/10
- Required Medical Examination by Dr. 09/09/09
- Health Care Provider Detail by Dr. 08/19/09
- PEER Review by Dr.— 10/08/08
- Office visits notes by Dr. 10/23/09 to 01/18/10

PATIENT CLINICAL HISTORY [SUMMARY]:

This patient sustained a work related injury on xx/xx/xx when she was moving furniture and experienced the onset of low back pain. She has been diagnosed with internal disc disruption and spinal stenosis at L5-S1. The patient has been treated with physical therapy and medications. The note for Dr. states that the patient's medications are working well and has been prescribed Flexeril 10mg 3 times a day, Valium 10mg for sleep, Norco 10/325 4 times a day and Xanax 3mg at bedtime. The patient has had refill prescriptions for these medications that have been denied payment twice.

ANALYSIS AND EXPLANATION OF THE DECISION INCLUDE CLINICAL BASIS, FINDINGS AND CONCLUSIONS USED TO SUPPORT THE DECISION.

This patient sustained a lumbar strain superimposed on degenerative changes and she has developed chronic low back pain. Muscle relaxants (Flexeril) are indicated for the treatment of acute painful musculoskeletal conditions. No studies have demonstrated efficacy with their use for chronic low back pain and therefore, the use of this medication in this capacity is not medically indicated. Narcotic analgesics (Norco) should be prescribed for fixed periods of time. Long-term treatment with short-acting opiates is not effective or indicated treatment of chronic back pain. Therefore the use of this medication is not medically indicated. Benzodiazepines (Valium, Xanax) are not indicated for the treatment of chronic low back pain. Therefore, the use of these medications is not medically indicated.

A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR OTHER CLINICAL BASIS USED TO MAKE THE DECISION:

ACOEM- AMERICAN COLLEGE OF OCCUPATIONAL & ENVIRONMENTAL MEDICINE UM KNOWLEDGEBASE	
AHCPR- AGENCY FOR HEALTHCARE RESEARCH & QUALI GUIDELINES	ΤY

☐ DWC- DIVISION OF WORKERS COMPENSATION POLICIES OR GUIDELINES
☐ EUROPEAN GUIDELINES FOR MANAGEMENT OF CHRONIC LOW BACK PAIN
☐ INTERQUAL CRITERIA
MEDICAL JUDGEMENT, CLINICAL EXPERIENCE AND EXPERTISE IN ACCORDANCE WITH ACCEPTED MEDICAL STANDARDS
☐ MERCY CENTER CONSENSUS CONFERENCE GUIDELINES
☐ MILLIMAN CARE GUIDELINES
☑ ODG- OFFICIAL DISABILITY GUIDELINES & TREATMENT GUIDELINES
☐ PRESSLEY REED, THE MEDICAL DISABILITY ADVISOR
☐ TEXAS GUIDELINES FOR CHIROPRACTIC QUALITY ASSURANCE & PRACTICE PARAMETERS
☐ TEXAS TACADA GUIDELINES
☐ TMF SCREENING CRITERIA MANUAL
☐ PEER REVIEWED NATIONALLY ACCEPTED MEDICAL LITERATURE (PROVIDE A DESCRIPTION)
OTHER EVIDENCE BASED, SCIENTIFICALLY VALID, OUTCOME FOCUSED GUIDELINES (PROVIDE A DESCRIPTION)